

Generalized Cluster Complexes via Quiver Representations

Bin Zhu¹

Department of Mathematical Sciences
Tsinghua University
100084 Beijing, P. R. China
E-mail: bzhu@math.tsinghua.edu.cn

February 2, 2008

Abstract

We give a quiver representation theoretic interpretation of generalized cluster complexes defined by Fomin and Reading. By using d -cluster categories which are defined by Keller as triangulated orbit categories of (bounded) derived categories of representations of valued quivers, we define a d -compatibility degree $(-||-)$ on any pair of “colored” almost positive real Schur roots which generalizes previous definitions on the non-colored case, and call two such roots compatible provided the d -compatibility degree of them is zero. Associated to the root system Φ corresponding to the valued quiver, by using this compatibility relation, we define a simplicial complex which has colored almost positive real Schur roots as vertices and d -compatible subsets as simplices. If the valued quiver is an alternating quiver of a Dynkin diagram, then this complex is the generalized cluster complex defined by Fomin and Reading.

Key words. Colored almost positive real Schur root, generalized cluster complex, d -cluster category, d -cluster tilting object, d -compatibility degree.

Mathematics Subject Classification. 05A15, 16G20, 16G70. 17B20.

1 Introduction

Generalized cluster complexes associated to finite root systems are introduced by Fomin and Reading [FR2]. They have some nice properties, see [AT] and the references there. They are a generalization of cluster complexes (so-called generalized associahedra) associated to the same root systems introduced in [FZ2, FZ3]. Cluster complexes describe the combinatorial structure of cluster algebras which were introduced by Fomin-Zelevinsky [FZ1] in order to give an algebraic and combinatorial framework for the canonical basis, see [FR1] for a nice survey on this combinatorics and also cluster combinatorics of root systems. In [MRZ], Marsh-Reineke-Zelevinsky use “decorated” quiver representations and tilting theory to give a quiver interpretation of cluster complexes. This connection between tilting theory and cluster combinatorics leads Buan-Marsh-Reineke-Reiten-Todorov [BMRRT] to introduce cluster categories for a categorical model for cluster algebras, see also [CCS] for type A_n . Cluster categories are the orbit categories $\mathcal{D}/\tau^{-1}[1]$ of derived categories of hereditary categories arising from the action of subgroup $\langle \tau^{-1}[1] \rangle$ of

¹Supported by the NSF of China (Grants 10471071) and by the Leverhulme Trust through the network ‘Algebras, Representations and Applications’.

the automorphism group. They are triangulated categories [Ke] and now they have become a successful model for acyclic cluster algebras [BMR, CC, CK], see also the surveys [BM, Rin] and the references there for recent developments and background of cluster tilting theory.

d -cluster categories $\mathcal{D}/\tau^{-1}[d]$ as a generalization of cluster categories, were introduced by Keller [Ke], Thomas [Th], for $d \in \mathbf{N}$. They are studied by Keller and Reiten [KR], Y. Palu [Pa], [ABST], see also [BaM] for a geometric description of d -cluster categories of type A_n . d -cluster categories are triangulated categories with Calabi-Yau dimension $d+1$. When $d=1$, the cluster categories are recovered.

The aim of this paper is to give, not only a quiver representation theoretic interpretation of all key ingredients in defining generalized cluster complexes using d -cluster categories, but also a generalization of generalized cluster complexes to infinite root systems (compare Remark 3.13 in [FR2], there the authors asked whether there was such an extension). For simply-laced Dynkin case, Thomas [Th] gives a realization of generalized cluster complexes by defining the d -cluster categories.

The paper is organized as follows: In the first two parts, we recall the well-known facts on d -cluster categories and (generalized) cluster complexes of finite root systems. In particular, we recall and generalize the BGP-reflection functors for cluster categories [Z1, Z2] to d -cluster categories. In the third part, we prove some properties of d -cluster tilting objects, including that any basic d -cluster tilting object contains exactly n indecomposable direct summands. In the final section, for any root system Φ , using a d -cluster category $\mathcal{C}_d(\mathcal{H})$, we define a d -compatibility degree on any pair of colored almost positive real Schur roots. Using the d -compatibility degree, we define a generalized cluster complex associated to Φ , which has colored almost positive real Schur roots as the vertices, and any subset forms a face if and only if any two elements of this subset are d -compatible. This simplicial complex is isomorphic to the cluster complex of d -cluster category $\mathcal{C}_d(\mathcal{H})$. If Φ is a finite root system, and if we take $\mathcal{H}_0$ to be the category of representations of an alternating quiver corresponding to Φ , then our generalized cluster complex is the usual generalized cluster complex $\Delta^d(\Phi)$ defined by Fomin and Reading in [FR2].

2 Basics on d -cluster categories

In this section, we collect some basic materials and fix notation which we will use later on.

A valued graph $(\Gamma, \mathbf{d})$ is a finite set of vertices $1, \dots, n$, together with non-negative integers d_{ij} for all pairs $i, j \in \Gamma$ such that $d_{ii} = 0$ and there exist positive integers $\{\varepsilon_i\}_{i \in \Gamma}$ satisfying

$$d_{ij}\varepsilon_j = d_{ji}\varepsilon_i, \text{ for all } i, j \in \Gamma.$$

A pair $\{i, j\}$ of vertices is called an edge of $(\Gamma, \mathbf{d})$ if $d_{ij} \neq 0$. An orientation Ω of a valued graph $(\Gamma, \mathbf{d})$ is given by prescribing for each edge $\{i, j\}$ of $(\Gamma, \mathbf{d})$ an order (indicated by an arrow $i \rightarrow j$). For simplicity, we denote a valued graph by Γ , and a valued quiver by (Γ, Ω) .

Let (Γ, Ω) be a valued quiver. We always assume that the valued quiver (Γ, Ω) contains no oriented cycles. Such orientation Ω is called admissible. Let K be a field and $\mathbf{M} =$

$(F_i, {}_iM_j)_{i,j \in \Gamma}$ a reduced K -species of (Γ, Ω) ; that is, for all $i, j \in \Gamma$, ${}_iM_j$ is an $F_i - F_j$ -bimodule, where F_i and F_j are division rings which are finite dimensional vector spaces over K and $\dim({}_iM_j)_{F_j} = d_{ij}$ and $\dim_K F_i = \varepsilon_i$. We denote by $\mathcal{H}$ the category of finite dimensional representations of $(\Gamma, \Omega, \mathcal{M})$. It is a hereditary abelian category [DR]. Let Φ be the root system of the Kac-Moody Lie algebra corresponding to the graph Γ . We assume that $P_1, \dots, P_n$ are the non-isomorphic indecomposable projective representations in $\mathcal{H}$, $E_1, \dots, E_n$ are the simple representations with dimension vectors $\alpha_1, \dots, \alpha_n$, and $\alpha_1, \dots, \alpha_n$ are the simple roots in Φ . We use $D(-)$ to denote $\text{Hom}_K(-, K)$ which is a duality of $\mathcal{H}$.

Denote by $\mathcal{D} = D^b(\mathcal{H})$ the bounded derived category of $\mathcal{H}$ with shift functor [1].

2.1 d -cluster categories

The derived category $\mathcal{D}$ has Auslander-Reiten triangles, and the Auslander-Reiten translate τ is an automorphism of $\mathcal{D}$. Fix a positive integer d , and denote by $F_d = \tau^{-1}[d]$, it is an automorphism of $\mathcal{D}$. The d -cluster category of $\mathcal{H}$ is defined in [Ke, Th]:

We denote by $\mathcal{D}/F_d$ the corresponding factor category. The objects are by definition the F_d -orbits of objects in $\mathcal{D}$, and the morphisms are given by

$$\text{Hom}_{\mathcal{D}/F_d}(\tilde{X}, \tilde{Y}) = \oplus_{i \in \mathbf{Z}} \text{Hom}_{\mathcal{D}}(X, F_d^i Y).$$

Here X and Y are objects in $\mathcal{D}$, and $\tilde{X}$ and $\tilde{Y}$ are the corresponding objects in $\mathcal{D}/F_d$ (although we shall sometimes write such objects simply as X and Y).

Definition 2.1. [Ke, Th] *The orbit category $\mathcal{D}/F_d$ is called the d -cluster category of $\mathcal{H}$ (or of (Γ, Ω)), which is denoted by $\mathcal{C}_d(\mathcal{H})$, sometimes denoted by $\mathcal{C}_d(\Omega)$.*

By [Ke], the d -cluster category is a triangulated category with shift functor [1] which is induced by shift functor in $\mathcal{D}$, the projection $\pi : \mathcal{D} \longrightarrow \mathcal{D}/F$ is a triangle functor. When $d = 1$, this orbit category is called the cluster category of $\mathcal{H}$, denoted by $\mathcal{C}(\mathcal{H})$ (sometimes denoted by $\mathcal{C}(\Omega)$).

$\mathcal{H}$ is a full subcategory of $\mathcal{D}$ consisting of complexes concentrated in degree 0, then passing to $\mathcal{C}_d(\mathcal{H})$ by the projection π , $\mathcal{H}$ is a (possibly not full) subcategory of $\mathcal{C}_d(\mathcal{H})$. For any $i \in \mathbf{Z}$, we use $(\mathcal{H})[i]$ to denote the copy of $\mathcal{H}$ under the i -th shift $[i]$ as a subcategory of $\mathcal{C}_d(\mathcal{H})$. In this way, we have that $(\text{ind } \mathcal{H})[i] = \{M[i] \mid M \in \text{ind } \mathcal{H}\}$. For any object M in $\mathcal{C}_d(\mathcal{H})$, $\text{add } M$ denotes the full subcategory of $\mathcal{C}_d(\mathcal{H})$ consisting of direct summands of direct sums of copies of M .

For $X, Y \in \mathcal{C}_d(\mathcal{H})$, we will use $\text{Hom}(X, Y)$ to denote the Hom-space $\text{Hom}_{\mathcal{C}_d(\mathcal{H})}(X, Y)$ in the d -cluster category $\mathcal{C}_d(\mathcal{H})$ throughout the paper. Define $\text{Ext}^i(X, Y)$ to be $\text{Hom}(X, Y[i])$.

We summarize some known facts about d -cluster categories [BMRRT, Ke].

Proposition 2.2. 1. $\mathcal{C}_d(\mathcal{H})$ has Auslander-Reiten triangles and Serre functor $\Sigma = \tau[1]$, where τ is the AR-translate in $\mathcal{C}_d(\mathcal{H})$, which is induced from AR-translate in $\mathcal{D}$.

2. $\mathcal{C}_d(\mathcal{H})$ is a Calabi-Yau category of CY-dimension $d + 1$.

3. $\mathcal{C}_d(\mathcal{H})$ is a Krull-Remark-Schmidt category.
4. $\text{ind}\mathcal{C}_d(\mathcal{H}) = \bigcup_{i=0}^{d-1} (\text{ind}\mathcal{H})[i] \cup \{P_j[d] \mid 1 \leq j \leq n\}$.

Proof. 1. This is Proposition 1.3 [BMRRT] and Corollary 1 in Section 8.4 of [Ke].

2. It is proved in Corollary 1 in Section 8.4 of [Ke]

3. This is proved in Proposition 1.2 [BMRRT]

4. The proof for $d = 1$ is given in Proposition 1.6 [BMRRT], which can be modified for the general d .

□

From Proposition 2.2, we define the degree for every indecomposable object in $\mathcal{C}_d(\mathcal{H})$ as follows:

Definition 2.3. For any indecomposable object $X \in \mathcal{C}_d(\mathcal{H})$, we call the non-negative integer $\min\{k \in \mathbf{Z}_{\geq 0} \mid X \cong M[k] \text{ in } \mathcal{C}_d(\mathcal{H}), \text{ for some } M \in \text{ind}\mathcal{H}\}$ the degree of X , denoted by $\deg X$.

From Definition 2.3, any indecomposable object X of degree k is isomorphic to $M[k]$ in $\mathcal{C}_d(\mathcal{H})$, where M is an indecomposable representation in $\mathcal{H}$; $0 \leq \deg X \leq d$, X has degree d if and only if $X \cong P[d]$ in $\mathcal{C}_d(\mathcal{H})$ for some indecomposable projective object $P \in \mathcal{H}$; and X has degree 0 if and only if $X \cong M[0]$ in $\mathcal{C}_d(\mathcal{H})$ for some indecomposable object $M \in \mathcal{H}$. Here $M[0]$ means regarding the object M of $\mathcal{H}$ as a complex concentrated in degree 0.

2.2 BGP-reflection functors

If T is a tilting object in $\mathcal{H}$, then the endomorphism algebra $A = \text{End}_{\mathcal{H}}(T)$ is called a tilted algebra. The tilting functor $\text{Hom}_{\mathcal{H}}(T, -)$ induces an equivalence $\text{RHom}(T, -) : D^b(\mathcal{H}) \rightarrow D^b(A)$, where $\text{RHom}(T, -)$ is the derived functor of $\text{Hom}_{\mathcal{H}}(T, -)$.

Any standard triangle functor $G : D^b(\mathcal{H}) \rightarrow D^b(\mathcal{H}')$ induces a well-defined functor $\tilde{G} : \mathcal{C}_d(\mathcal{H}) \rightarrow \mathcal{C}_d(\mathcal{H}')$ with the following commutative diagram [Ke, Z1]:

$$\begin{array}{ccc} D^b(\mathcal{H}) & \xrightarrow{G} & D^b(\mathcal{H}') \\ \downarrow & & \downarrow \\ \mathcal{C}_d(\mathcal{H}) & \xrightarrow{\tilde{G}} & \mathcal{C}_d(\mathcal{H}') \end{array}$$

The following result is proved in [Z1, Z2].

Proposition 2.4. If $G : D^b(\mathcal{H}) \rightarrow D^b(\mathcal{H}')$ is a triangle equivalence, then $\tilde{G}$ is also an equivalence of triangulated categories.

Let k be a vertex in the valued quiver (Γ, Ω) , the reflection of (Γ, Ω) at k is the valued quiver $(\Gamma, s_k\Omega)$, where $s_k\Omega$ is the orientation of Γ obtained from Ω by reversing all arrows starting or ending at k . The corresponding category of representations of $(\Gamma, s_k\Omega, \mathcal{M})$ is denoted simply by $s_k\mathcal{H}$. If k is a sink in the valued quiver (Γ, Ω) , then k is a source of

$(\Gamma, s_k\Omega)$, and the reflection of $(\Gamma, s_k\Omega)$ at k is (Γ, Ω) . Let k be a sink in (Γ, Ω) . Then P_k is a simple projective representation and $T = \bigoplus_{j \neq k} P_j \oplus \tau^{-1}P_k$ is a tilting representation in $\mathcal{H}$ [Rin]. The tilting functor $S_k^+ = \text{Hom}_{\mathcal{H}}(T, -)$ is a so-called BGP-reflection functor, and its derived functor $\text{RHom}(T, -)$ is a triangle equivalence from $D^b(\mathcal{H})$ to $D^b(s_k\mathcal{H})$, which is also denoted by S_k^+ . Similarly, one has BGP-reflection functors S_k^- for sources k .

Definition 2.5. *The induced functors $\widetilde{S}_k^+ : \mathcal{C}_d(\mathcal{H}) \longrightarrow \mathcal{C}_d(s_k\mathcal{H})$ for sinks k and $\widetilde{S}_k^- : \mathcal{C}_d(\mathcal{H}) \longrightarrow \mathcal{C}_d(s_k\mathcal{H})$ for sources k are called BGP-reflection functors of d -cluster categories.*

Remark 2.6. *When $d = 1$, BGP-reflection functors are discussed in [Z1].*

We remind the reader that $\mathcal{H}$ (or $\mathcal{H}'$) is the category of representations of the valued quiver (Γ, Ω) ($(\Gamma, s_k\Omega)$, respectively); the P_i (respectively, the P'_i) are the indecomposable projective representations in $\mathcal{H}$ (respectively, $\mathcal{H}'$) and the E_i (respectively, the E'_i) are the corresponding simple representations which are the tops of the P_i (respectively, the P'_i), for $i = 1, \dots, n$.

We recall from Proposition 2.2 and Definition 2.3 that any indecomposable object Y in $\mathcal{C}_d(\mathcal{H})$ is isomorphic to $X[i]$ where $X \in \text{ind}\mathcal{H}$ and i is the degree of Y . Keeping this notation, we have the following proposition which gives the images of indecomposable objects in $\mathcal{C}_d(\mathcal{H})$ under the BGP-reflection functor $\widetilde{S}_k^+$.

Proposition 2.7. *Let k be a sink of the valued quiver (Γ, Ω) and Y an indecomposable object in $\mathcal{C}_d(\mathcal{H})$ with degree i . Then $Y \cong X[i]$ for an indecomposable representation X in $\mathcal{H}$, and*

$$\widetilde{S}_k^+(X[i]) = \begin{cases} P'_k[d] & \text{if } X \cong P_k (\cong E_k) \text{ and } i = 0 \\ E'_k[i-1] & \text{if } X \cong P_k (\cong E_k) \text{ and } 0 < i \leq d \\ P'_j[d] & \text{if } X \cong P_j \not\cong P_k \text{ and } i = d \\ S_k^+(X)[i] & \text{otherwise;} \end{cases}$$

Proof. The statement in the proposition was proved in [Z1, Z2] when $d = 1$. The proof for the case $d > 1$ is the same as there. We give a sketch of the proof for the convenience of readers. The BGP-reflection functor $S_k^+ : \mathcal{H} \longrightarrow s_k\mathcal{H}$ induces a triangle equivalence $D^b(\mathcal{H}) \longrightarrow D^b(s_k\mathcal{H})$, denoted also by S_k^+ . It induces an equivalence $\text{ind}D^b(\mathcal{H}) \longrightarrow \text{ind}D^b(s_k\mathcal{H})$. For any indecomposable object $X[i] \in \text{ind}D^b(\mathcal{H})$, it is not hard to show that $S_k^+(X[i]) = S_k^+(X)[i]$ for $X \not\cong P_k$ (note that $P_k = E_k$ since k is a sink in (Γ, Ω)), and $S_k^+(P_k[i]) = E'_k[i-1]$ for $i \in \mathbf{Z}$ (compare [Z1] or [Z2]). Since E'_k is an injective representation in $s_k\mathcal{H}$, we have $\tau P'_k[i] = E'_k[i-1]$ in $D^b(s_k\mathcal{H})$. Now passing to the d -cluster category $\mathcal{C}_d(\mathcal{H})$ (which is an orbit category of the derived category $D^b(\mathcal{H})$), we get the images of indecomposable objects of $\mathcal{C}_d(\mathcal{H})$ under $\widetilde{S}_k^+$ as stated in the proposition. $\square$

3 Cluster combinatorics of root systems

For a valued graph Γ , we denote by $\Phi = \Phi^+ \cup \Phi^-$ the set of roots of the corresponding Kac-Moody Lie algebra.

Definition 3.1. 1. The set of almost positive roots is

$$\Phi_{\geq -1} = \Phi^+ \cup \{-\alpha_i \mid i = 1, \dots, n\}.$$

2. Denote by $\Phi_{\geq -1}^{re}$ the subset of $\Phi_{\geq -1}$ consisting of the positive real roots together with the negatives of the simple roots.

When Φ is of finite type, $\Phi_{\geq -1} = \Phi_{\geq -1}^{re}$.

Definition 3.2. Let s_i be the Coxeter generator of the Weyl group of Φ corresponding to $i \in \Gamma_0$. We call the following map the "truncated simple reflection" σ_i of $\Phi_{\geq -1}$ [FZ2]:

$$\sigma_i(\alpha) = \begin{cases} \alpha & \alpha = -\alpha_j, j \neq i \\ s_i(\alpha) & \text{otherwise.} \end{cases}$$

It is easy to see that σ_i is an automorphism of $\Phi_{\geq -1}^{re}$.

3.1 Cluster complexes of finite root systems

In this first paragraph, we don't assume that Γ is a Dynkin diagram (i.e. of finite type). Let $i_1, \dots, i_n$ be an admissible ordering of Γ with respect to Ω , i.e. i_t is a sink with respect to $s_{i_{t-1}} \cdots s_{i_2} s_{i_1} \Omega$ for any $1 \leq t \leq n$. Denote by $R_\Omega = \sigma_{i_n} \cdots \sigma_{i_1}$. This is an automorphism of $\Phi_{\geq -1}$ and does not depend on the choice of admissible ordering of Γ with respect to Ω . It is the automorphism induced by Auslander-Reiten translation τ in $\mathcal{C}(\mathcal{H})$ (compare [Z1, Z2]).

In the rest of this subsection, we always assume that Γ is a valued Dynkin graph, which is not necessarily connected. Fomin and Zelevinsky [FZ2] associate a nonnegative integer $(\alpha || \beta)$, known as the compatibility degree, to each pair α, β of almost positive roots. This is defined in the following way: Let Ω_0 denote one of the alternating orientations of Γ , and Γ^+ (respectively, Γ^-) the set of sinks (respectively, sources) of (Γ, Ω_0) respectively. Define:

$$\tau_\pm = \prod_{i \in \Gamma^\pm} \sigma_i.$$

Then $R_{\Omega_0} = \tau_- \tau_+$, which is simply denoted by R .

Denote by $n_i(\beta)$ the coefficient of α_i in the expansion of β in terms of the simple roots $\alpha_1, \dots, \alpha_n$. Then $(||)$ is uniquely defined by the following two properties:

$$\begin{aligned} (*) \quad & (-\alpha_i || \beta) = \max([\beta : \alpha_i], 0), \\ (**) \quad & (\tau_\pm \alpha || \tau_\pm \beta) = (\alpha || \beta), \end{aligned}$$

for any $\alpha, \beta \in \Phi_{\geq -1}$, any $i \in \Gamma$.

Two almost positive roots α, β are called compatible if $(\alpha || \beta) = 0$.

The cluster complex $\Delta(\Phi)$ associated to the finite root system Φ is defined in [FZ2].

Definition 3.3. *The cluster complex $\Delta(\Phi)$ is a simplicial complex on the ground set $\Phi_{\geq -1}$. Its faces are mutually compatible subsets of $\Phi_{\geq -1}$. The facets of $\Delta(\Phi)$ are called the (root-)clusters associated to Φ .*

3.2 Generalized cluster complexes of finite root systems

At the beginning of this subsection, we assume that Γ is arbitrary valued graph, which is not necessarily connected, except where we express specifically. As before Φ denotes the set of roots of the corresponding Lie algebra, and $\Phi_{\geq -1}$ denotes the set of almost positive roots. Fix a positive integer d , for any $\alpha \in \Phi^+$, following [FR2], we call $\alpha^1, \dots, \alpha^d$ the d “colored” copies of α .

Definition 3.4. *[FR2] The set of colored almost positive roots is*

$$\Phi_{\geq -1}^d = \{\alpha^i : \alpha \in \Phi_{>0}, i \in \{1, \dots, d\}\} \bigcup \{(-\alpha_i)^1 : 1 \leq i \leq n\}.$$

When Γ is a Dynkin graph, the root system Φ of the corresponding Lie algebra is finite. In this case the generalized cluster complex $\Delta^d(\Phi)$ is defined on the ground set $\Phi_{\geq -1}^d$ and using the binary compatibility relation on $\Phi_{\geq -1}^d$. This binary compatibility relation is a natural generalization of binary compatibility relation on $\Phi_{\geq -1}$ which we now recall from [FR2].

For a root $\beta \in \Phi_{\geq -1}$, let $t(\beta)$ denote the smallest t such that $R^t(\beta)$ is a negative root.

Definition 3.5. *[FR2] Two colored roots $\alpha^k, \beta^l \in \Phi_{\geq -1}^d$ are called compatible if and only if one of the following conditions is satisfied:*

1. $k > l$. $t(\alpha) \leq t(\beta)$, and the roots $R(\alpha)$ and β are compatible (in the original “non-colored” sense)
2. $k < l$. $t(\alpha) \geq t(\beta)$, and the roots α and $R(\beta)$ are compatible
3. $k > l$. $t(\alpha) > t(\beta)$, and the roots α and β are compatible
4. $k < l$. $t(\alpha) < t(\beta)$, and the roots α and β are compatible
5. $k = l$. and the roots α and β are compatible

Now we are ready to recall the definition of generalized cluster complex $\Delta^d(\Phi)$ for a finite root system Φ .

Definition 3.6. *[FR2] $\Delta^d(\Phi)$ has $\Phi_{\geq -1}^d$ as the set of vertices, its simplices are mutually compatible subsets of $\Phi_{\geq -1}^d$. The subcomplex of $\Delta^d(\Phi)$ which has $\Phi_{>0}^d$ as the set of vertices is denoted by $\Delta_+^d(\Phi)$*

Now we generalize the definition of R_d [FR2] for finite root system to an arbitrary root system.

Definition 3.7. *Let (Γ, Ω) be a valued quiver. For $\alpha^k \in \Phi_{\geq -1}^d$, we set*

$$R_{d,\Omega}(\alpha^k) = \begin{cases} \alpha^{k+1} & \text{if } \alpha \in \Phi_{>0} \text{ and } k < d; \\ (R_\Omega(\alpha))^1 & \text{otherwise,} \end{cases}$$

Remark 3.8. If (Γ, Ω_0) is a valued Dynkin graph with an alternating orientation, then the automorphism R of $\Phi_{\geq -1}$ defined by Fomin-Zelevinsky [FZ2] is R_{Ω_0} , hence R_{d, Ω_0} is the usual one (R_d) defined by Fomin-Reading [FR2].

Theorem 3.9. [FR2] Let Φ be a finite root system. The compatibility relation on $\Phi_{\geq -1}^d$ has the following properties:

1. α^k is compatible with β^l if and only if $R_d(\alpha^k)$ is compatible with $R_d(\beta^l)$.
2. $(-\alpha_i)^1$ is compatible with β^l if and only if $n_i(\beta) = 0$.

Furthermore, conditions 1 – 2 uniquely determine this relation.

Now we generalize the “truncated simple reflections” of $\Phi_{\geq -1}$ to the colored almost positive roots. Let Φ be an arbitrary root system (not necessarily of finite type).

Definition 3.10. Let s_k be the Coxeter generator of the Weyl group of Φ corresponding to $k \in \Gamma_0$. We define the following map $\sigma_{k,d}$ of $\Phi_{\geq -1}^d$:

$$\sigma_{k,d}(\alpha^i) = \begin{cases} \alpha_k^d & \text{if } i = 1, \text{ and } \alpha = -\alpha_k, \\ \alpha_k^{i-1} & \text{if } 1 < i \leq d, \text{ and } \alpha = \alpha_k, \\ (-\alpha_j)^1 & \text{if } i = 1, \text{ and } \alpha = -\alpha_j, \ j \neq k \\ (s_k(\alpha))^i & \text{otherwise.} \end{cases}$$

$\sigma_{k,d}$ is a bijection of $\Phi_{\geq -1}^d$. We call it a d -truncated simple reflection of $\Phi_{\geq -1}^d$.

4 d -cluster tilting in d -cluster categories

Let $\mathcal{C}_d(\mathcal{H})$ be a d -cluster category of type $\mathcal{H}$, where $\mathcal{H}$ is the category of representations of the valued quiver (Γ, Ω) . It is a Calabi-Yau triangulated category with CY-dimension $d + 1$.

Definition 4.1. 1. An object X in $\mathcal{C}_d(\mathcal{H})$ is called exceptional if $\text{Ext}^i(X, X) = 0$, for any $1 \leq i \leq d$.

2. An object X is called a d -cluster tilting object if it satisfies the property: $Y \in \text{add} X$ if and only if $\text{Ext}^i(X, Y) = 0$ for $1 \leq i \leq d$.

3. An object X is called almost complete tilting if there is an indecomposable object Y such that $X \oplus Y$ is a d -cluster tilting object. Such an indecomposable object Y is called a complement of X .

Proposition 4.2. 1. For an object X in $\mathcal{H}$, X is exceptional in $\mathcal{H}$ i.e. $\text{Ext}_{\mathcal{H}}^1(X, X) = 0$ if and only if $X[0]$ is exceptional in $\mathcal{C}_d(\mathcal{H})$.

2. Any indecomposable exceptional object X in $\mathcal{C}_d(\mathcal{H})$ is of the form $M[i]$ with M being an exceptional representation in $\mathcal{H}$ and $0 \leq i \leq d - 1$ or of the form $P_j[d]$ for some $1 \leq j \leq n$. In particular, if Γ is a Dynkin graph, then any indecomposable object in $\mathcal{C}_d(\mathcal{H})$ is exceptional.

3. Suppose $d > 1$. Then $\text{End}_{\mathcal{C}_d(\mathcal{H})} X$ is a division algebra for any indecomposable exceptional object X .
4. Suppose $d > 1$. Let P be a projective representation in $\mathcal{H}$ and X a representation in $\mathcal{H}$. Then for any $-d \leq i \leq d$, $\text{Ext}^1(P, X[i]) = 0$ except possibly for $i \in \{-1, d-1, d\}$.

Proof. 1. Let $X \in \mathcal{H}$ be exceptional. We will prove that $\text{Ext}^i(X, X) = 0$, for any $i \in \{1, \dots, d\}$. By definition, we have that $\text{Ext}^i(X, X) = \oplus_{k \in \mathbf{Z}} \text{Ext}_{\mathcal{D}}^i(X, \tau^{-k} X[kd]) = \text{Ext}_{\mathcal{D}}^i(X, X) \oplus \text{Ext}_{\mathcal{D}}^i(X, \tau X[-d])$. In this sum, the first summand $\text{Ext}_{\mathcal{D}}^i(X, X) = 0, \forall i \geq 1$, while the second summand $\text{Ext}_{\mathcal{D}}^i(X, \tau X[-d]) \cong \text{Hom}_{\mathcal{D}}(X, \tau X[i-d])$, which is zero when $i < d$, and is isomorphic to $\text{Ext}_{\mathcal{D}}^1(X, X) = 0$ when $i = d$. This proves that X is exceptional in $\mathcal{C}_d(\mathcal{H})$. The proof for the converse follows directly from definition.

2. The statements follow from Proposition 2.2(4) and Definition 4.1, also using part 1 and the fact that the shift is an autoequivalence.
3. Let X be an indecomposable exceptional representation in $\mathcal{H}$ and suppose $d > 1$. It follows from the definition of the orbit category that $\text{End}_{\mathcal{C}_d(H)} X \cong \oplus_{m \in \mathbf{Z}} \text{Hom}_{\mathcal{D}}(X, \tau^{-m} X[dm]) \cong \text{End}_{\mathcal{H}} X$. The last isomorphism holds due to the facts: $\text{Hom}_{\mathcal{D}}(X, \tau^m X[-md]) \cong \text{Hom}_{\mathcal{D}}(X[md], \tau^m X) \cong \text{Ext}_{\mathcal{D}}^1(\tau^{m-1} X, X[md]) = 0$ for any positive integer m ; and $\text{Hom}_{\mathcal{D}}(X, \tau^{-m} X[md]) \cong \text{Hom}_{\mathcal{D}}(\tau^m X, X[md])$, which is also zero since $md > 1$ (we use the assumption $d > 1$ here) for any positive integer m . Then $\text{End}_{\mathcal{C}_d(H)} X$ is a division algebra since $\text{End}_{\mathcal{H}} X$ is a division algebra. Since any indecomposable exceptional object M in $\mathcal{C}_d(H)$ is some shift $X[i]$ of an indecomposable exceptional representation X in $\mathcal{H}$, $\text{End}_{\mathcal{C}_d(H)} M = \text{End}_{\mathcal{C}_d(H)} X[i] \cong \text{End}_{\mathcal{C}_d(H)} X$ is a division algebra.
4. Suppose $d > 1$. Let P be a projective representation in $\mathcal{H}$ and X a representation in $\mathcal{H}$. Then for any $-d \leq i \leq d$, $\text{Ext}^1(P, X[i]) = \oplus_{k \in \mathbf{Z}} \text{Ext}_{\mathcal{D}}^1(P, \tau^{-k} X[dk+i]) \cong \text{Ext}_{\mathcal{D}}^1(P, \tau X[-d+i]) \oplus \text{Ext}_{\mathcal{D}}^1(P, X[i])$. Now if $i \neq -1, d-1, d$, then $\text{Ext}_{\mathcal{D}}^1(P, \tau X[-d+i]) = 0 = \text{Ext}_{\mathcal{D}}^1(P, X[i])$. Then for any $-d \leq i \leq d$, $\text{Ext}^1(P, X[i]) = 0$ except for $i = -1, d-1$ and d .

□

Remark 4.3. Any basic (i.e. multiplicity-free) exceptional object contains at most $(d+1)n$ non-isomorphic indecomposable direct summands.

Proof. Let X be a basic exceptional object in $\mathcal{C}_d(\mathcal{H})$. Then any indecomposable direct summand of X is exceptional, hence by Proposition 4.2 (2), we write M as $M = \oplus_{k=0}^{d-1} \oplus_{i \in I_k} M_{i,k}[k]$ with $M_{i,k}$ being an indecomposable exceptional representation. Therefore $\oplus_{i \in I_k} M_{i,k}$ is an exceptional object in hereditary category $\mathcal{H}$, hence the number of direct summands is at most n , i.e. $|I_k| \leq n$. Then the number of indecomposable direct summands of M is at most $(d+1)n$. □

For any pair of objects T, X in $\mathcal{C}_d(\mathcal{H})$, due to the Calabi-Yau property of $\mathcal{C}_d(\mathcal{H})$, we have that $\text{Ext}^i(X, T) = 0$ for $1 \leq i \leq d$ if and only if $\text{Ext}^i(T, X) = 0$ for $1 \leq i \leq d$. Hence from Remark 4.3 and Definition 4.1, T is a d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$ if and

only if $\text{add}T$ is a maximal d -orthogonal subcategory of $\mathcal{C}_d(\mathcal{H})$ in the sense of [I2]: i.e. $\text{add}T$ is contravariantly finite and covariantly finite in $\mathcal{C}_d(\mathcal{H})$, and satisfies the property: $X \in \text{add}T$ if and only if $\text{Ext}^i(X, T) = 0$ for $1 \leq i \leq d$ if and only if $\text{Ext}^i(T, X) = 0$ for $1 \leq i \leq d$. In the following we will prove that any basic d -cluster tilting object contains exactly n indecomposable direct summands. First of all, we recall some results from [I2] which hold in any $(d+1)$ -Calabi-Yau triangulated category.

Theorem 4.4. (*Iyama*) *Let X be an almost complete tilting object in $\mathcal{C}_d(\mathcal{H})$, and X_0 a complement of X . Then there are $d+1$ triangles:*

$$(*) \quad X_{i+1} \xrightarrow{g_i} B_i \xrightarrow{f_i} X_i \xrightarrow{\sigma_i} X_{i+1}[1]$$

where f_i is the minimal right $\text{add}X$ -approximation of X_i and g_i minimal left $\text{add}X$ -approximation of X_{i+1} , all X_i are indecomposable and complements of X , $i = 0, \dots, d$.

For the convenience of readers, we sketch the proof, for details, see [IY].

Proof. We suppose that $d > 1$, the same statement for $d = 1$ was proved in [BMRRT]. For the complement X_0 of X , we consider the minimal right $\text{add}X$ -approximation $f_0 : B_0 \rightarrow X_0$ of X_0 , extend f_0 to a triangle $X_1 \xrightarrow{g_0} B_0 \xrightarrow{f_0} X_0 \xrightarrow{\sigma_0} X_1[1]$. It is easy to see that X_1 is indecomposable, g_0 is the minimal left $\text{add}X$ -approximation of X_1 and $X \oplus X_1$ is an exceptional object in $\mathcal{C}_d(\mathcal{H})$ (compare [BMRRT]). It follows from Theorem 5.1 in [IY] that $X \oplus X_1$ is a d -cluster tilting object. Continuing this step, one can get complements $X_1, \dots, X_{d+1}$ with triangles $X_{i+1} \xrightarrow{g_i} B_i \xrightarrow{f_i} X_i \xrightarrow{\sigma_i} X_{i+1}[1]$ for $0 \leq i \leq d$, where f_i (g_i) is the minimal right (left, resp.) $\text{add}X$ -approximation of X_i (X_{i+1} , resp.) and $X \oplus X_i$ is a d -cluster tilting object. $\square$

Corollary 4.5. *With the same notation as Theorem 4.4, we have that $\sigma_d[d]\sigma_{d-1}[d-1] \cdots \sigma_1[1]\sigma_0 \neq 0$. In particular, $\text{Hom}(X_i, X_j[j-i]) \neq 0$ and $X_i \not\cong X_j, \forall 0 \leq i < j \leq d$.*

Proof. From Theorem 4.4, we have that $\sigma_0 \neq 0$ since the triangle $(*)$ at $i = 0$ in Theorem 4.4 is non-splitting. Suppose that $\sigma_d[d]\sigma_{d-1}[d-1] \cdots \sigma_1[1]\sigma_0 = 0$, then $\sigma_{d-1}[d-1] \cdots \sigma_1[1]\sigma_0 : X_0 \rightarrow X_d[d]$ factors through $f_d[d] : B_d[d] \rightarrow X_d[d]$ since we have a triangle $X_{d+1}[d] \xrightarrow{g_d[d]} B_d[d] \xrightarrow{f_d[d]} X_d[d] \xrightarrow{\sigma_d[d]} X_{d+1}[d+1]$. Since $\text{Hom}(X_0, B_d[d]) = \text{Ext}^d(X_0, B_d) = 0$, $\sigma_{d-1}[d-1] \cdots \sigma_1[1]\sigma_0 = 0$. Similarly $\sigma_{d-2}[d-2] \cdots \sigma_1[1]\sigma_0 = 0$, and finally, $\sigma_0 = 0$, a contradiction. Now we prove final statement: we have that $\sigma_{j-1}[j-1] \cdots \sigma_i \in \text{Hom}(X_i, X_j[j-i])$ and $\sigma_{j-1}[j-1] \cdots \sigma_i \neq 0$. Otherwise $\sigma_{j-1}[j-1] \cdots \sigma_i = 0$, and hence $\sigma_{d-1}[d-1] \cdots \sigma_1[1]\sigma_0 = 0$, a contradiction. Now suppose that $X_i \cong X_j$ for some $i < j$. Then $\text{Ext}^k(X_i, X_j) = 0$ for $1 \leq k \leq d$, a contradiction. Then $X_i \not\cong X_j$. $\square$

Now we state our main result of this section.

Theorem 4.6. *Any basic d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$ contains exactly n indecomposable direct summands.*

To prove the theorem, we need some technical lemmas.

Lemma 4.7. *Let $d > 1$ and $X = M[i]$, $Y = N[j]$ be indecomposable objects of degree i, j respectively in $\mathcal{C}_d(\mathcal{H})$. Suppose that $\text{Hom}(X, Y) \neq 0$. Then one of the following holds:*

- (1) *We have $i = j$ or $j - 1$ (provided $j \geq 1$);*
- (2) *We have $i = 0, i = d$ (and $M = P$) or $d - 1$ (provided $j = 0$).*

Proof. Let $d > 1$. Firstly we note that for any indecomposable object $X \in \mathcal{C}_d(\mathcal{H})$, $0 \leq \deg X \leq d$, $\deg X = d$ if and only if $X = P_i[d]$ for an indecomposable projective representation P_i . This implies that $-d \leq \deg Y - \deg X \leq d$ for indecomposable objects $X, Y \in \mathcal{C}_d(\mathcal{H})$. Let $X = M[i]$, $Y = N[j]$ be indecomposable objects of degree i, j respectively in $\mathcal{C}_d(\mathcal{H})$. We have $\text{Hom}(X, Y) \cong \text{Hom}(M, N[j - i]) = \oplus_{k \in \mathbf{Z}} \text{Hom}_{\mathcal{D}}(M, \tau^{-k} N[j - i + kd]) = \text{Hom}_{\mathcal{D}}(M, \tau N[j - i - d]) \oplus \text{Hom}_{\mathcal{D}}(M, N[j - i]) \oplus \text{Hom}_{\mathcal{D}}(M, \tau^{-1} N[j - i + d])$. The last equality holds due to $-d \leq j - i \leq d$, and $\text{Hom}_{\mathcal{D}}(M, \tau^{-k} N[j - i + kd]) = 0$ for $k \neq -1, 0, 1$. We divide the calculation of $\text{Hom}(X, Y)$ into three cases:

1. The case $-d < j - i < d$. We have that $\text{Hom}(X, Y) \cong \text{Hom}_{\mathcal{D}}(M, N[j - i]) \oplus \text{Hom}_{\mathcal{D}}(M, \tau^{-1} N[j - i + d])$. The first summand is zero when $j - i \neq 0, 1$ while the second is zero when $d + j - i \neq 1$ (equivalently $d + j - i > 1$ since $0 < d + j - i < 2d$).
2. The case $j - i = -d$. Then $j = 0, i = d$ ($M = P$). Then $\text{Hom}(X, Y) = \text{Hom}_{\mathcal{D}}(P, \tau^{-1} N)$.
3. The case $j - i = d$. Then $j = d$ ($N = P$), $i = 0$. Then $\text{Hom}(X, Y) = \text{Hom}_{\mathcal{D}}(M, \tau P) \oplus \text{Hom}_{\mathcal{D}}(M, P[d]) \oplus \text{Hom}_{\mathcal{D}}(M, \tau^{-1} P[2d]) = 0$.

Therefore if $\text{Hom}(X, Y) \neq 0$, then $\text{Hom}(M[0], N[j - i]) \neq 0$. Proof of (1). Suppose $j \geq 1$. Then combining with Case 3, we have that $-d < j - i < d$. We want to prove that if $j - i \neq 0, 1$ then $\text{Hom}(X, Y) = 0$ this will finish the proof of (1). Under the condition $j - i \neq 0, 1$, from Case 1, we have that $\text{Hom}(X, Y) \cong \text{Hom}_{\mathcal{D}}(M, \tau^{-1} N[j - i + d])$, which is zero for $d + j - i \neq 1$. But if $d + j - i = 1$, i.e. $i = d$, hence $M = P$ and $j = 1$. Then $\text{Hom}_{\mathcal{D}}(M, \tau^{-1} N[j - i + d]) = \text{Hom}_{\mathcal{D}}(P, \tau^{-1} N[1]) = 0$. We have finished the proof of (1). Proof of (2). Suppose $j = 0$. Then $-d \leq j - i \leq 0$. It follows from Cases 1-2 that $i = 0, i = d$ ($M = P$) or $i = d - 1$. This finishes the proof of (2). $\square$

Lemma 4.8. *If $d > 2$, then $\text{Ext}^2(M[i], N[i]) = 0$ for objects $M, N \in \mathcal{H}$, and any i .*

Proof. It is sufficient to prove that $\text{Ext}^2(M[0], N[0]) = 0$. From the definition of the orbit category $\mathcal{D}/\tau^{-1}[d]$, we have that

$$\text{Ext}^2(M[0], N[0]) = \text{Hom}(M[0], N[2]) = \oplus_{k \in \mathbf{Z}} \text{Hom}_{\mathcal{D}}(M, \tau^{-k} N[kd + 2]),$$

where each summand $\text{Hom}_{\mathcal{D}}(M, \tau^{-k} N[kd + 2])$ equals 0 since $kd + 2 \geq 2$ or $kd + 2 \leq -1$ by the condition $d > 2$. Hence $\text{Ext}^2(M[0], N[0]) = 0$. $\square$

Lemma 4.9. *Let $d > 1$ and $M, N \in \mathcal{H}$. Then $\text{Ext}^1(M[0], N[0]) \cong \text{Ext}_{\mathcal{H}}^1(M, N)$. Furthermore any non-split triangle between $M[0]$ and $N[0]$ in $\mathcal{C}_d(\mathcal{H})$ is induced from a non-split exact sequence between M and N in $\mathcal{H}$.*

Proof. Under the condition $d > 1$, it is easy to see that $\text{Ext}^1(M[0], N[0]) = \oplus_{k \in \mathbf{Z}} \text{Ext}_{\mathcal{D}}^1(M, \tau^{-k}N[2k]) = \text{Ext}_{\mathcal{D}}^1(M, N) = \text{Ext}_{\mathcal{H}}^1(M, N)$. This proves the first statement. Since $\mathcal{H} \subset \mathcal{C}_d(\mathcal{H})$ is a (not necessarily full) embedding, and any exact short sequence in $\mathcal{H}$ induces a triangle in $\mathcal{C}_d(\mathcal{H})$, the final statement then follows from the first statement. $\square$

Proof. (of Theorem 4.6): We assume that $d > 1$ since it was proved in [BMMRT] for $d = 1$. Let $M = \oplus_{i \in I} M_i[k_i]$ be a d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$, where all M_i are indecomposable representations in $\mathcal{H}$, $0 \leq k_i \leq d$ (when $k_i = d$, M_i is projective). One can assume that one of k_i is 0, otherwise one can replace M by a suitable shift of M . Denote by $\nu(M) = \max\{|k_i - k_j| \mid \forall i, j\}$. We prove $|I| = n$ by induction on $\nu(M)$, where $|I|$ denotes the cardinality of I . If $\nu(M) = 0$, i.e. $k_i = 0$ for all i , then $\oplus_{i \in I} M_i[0]$ is a d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$, hence a tilting object in $\mathcal{H}$. Then $|I| = n$. Now assume $\nu(M) = m > 0$. Without losing generality, we assume that $k_1 = \dots = k_t = m$ and $k_j < m$ for $j > t$. From the complement $X_0 = M_1[k_1]$ of $X = M \setminus M_1[k_1]$ (here we use $X \setminus X_1$ to denote a complement of X_1 in X for a direct summand X_1 of X), by Theorem 4.4, we have at least $d+1$ complements X_j , $j = 0, \dots, d$, which form the triangles $(*)$ in Theorem 4.4. In these triangles, it is easy to see that $f_i = 0$ if and only if $B_i = 0$ if and only if $g_i = 0$. We will prove that there are at least one of complements X_j with smaller degree than m . At first, we prove this statement for the special case that $m = 1$. We claim that the degree of X_1 is 0 or 1 in this case. Otherwise $X_1 = P[d]$ for some indecomposable projective representation P or $X_1 = Y[d-1]$ for some indecomposable representation Y . Write X_0 as $Z[1]$, where Z is an indecomposable representation in $\mathcal{H}$. If $X_1 = P[d]$, then $\text{Hom}(X_1, X_0[d]) = \text{Hom}(P[d], X_0[d]) \cong \text{Hom}(P, Z[1]) = 0$, this contradicts to the fact that $\text{Hom}(X_1, X_0[d]) \cong \text{Hom}(X_0, X_1[1])$ is not zero by Theorem 4.4 or Corollary 4.5. If $X_1 = Y[d-1]$, then X_1 has degree 1 when $d = 2$, and $\text{Hom}(X_1, X_0[d]) = \text{Hom}(Y[d-1], Z[d+1]) \cong \text{Ext}^2(Y, Z) = 0$ by Lemma 4.8. when $d > 2$, which also contradicts to the fact that $\text{Hom}(X_1, X_0[d]) \cong \text{Hom}(X_0, X_1[1])$ is not zero. This proves the statement that X_1 has degree 0 or 1. Now if there are no complements X_j of X with degree 0, then all X_j have degree 1. We prove that any three successive complements, say X_0, X_1, X_2 , can not have the same degree. If all degrees of $X_i, i = 0, 1, 2$, are the same, we can assume that all X_i have degree 0. By Lemma 4.9, we have non-split short exact sequences in $\mathcal{H}$:

$$\begin{aligned} 0 &\longrightarrow X_1 \longrightarrow B_0 \longrightarrow X_0 \longrightarrow 0 \\ 0 &\longrightarrow X_2 \longrightarrow B_1 \longrightarrow X_1 \longrightarrow 0 \end{aligned}$$

From the first short exact sequence, we have $\text{Ext}_{\mathcal{H}}^1(X_0, X_1) \not\cong 0$. By applying $\text{Hom}_{\mathcal{H}}(X_0, -)$ to the second exact sequence, we have the long exact sequence:

$$\dots \rightarrow \text{Ext}_{\mathcal{H}}^1(X_0, X_2) \rightarrow \text{Ext}_{\mathcal{H}}^1(X_0, B_1) \rightarrow \text{Ext}_{\mathcal{H}}^1(X_0, X_1) \rightarrow \text{Ext}_{\mathcal{H}}^2(X_0, X_2) \rightarrow \text{Ext}_{\mathcal{H}}^2(X_0, B_1).$$

Since $X \oplus X_0$ is a d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$ and $B_1 \in \text{add} X$, $\text{Ext}_{\mathcal{H}}^1(X_0, B_1) = 0$. Hence we have that $\text{Ext}_{\mathcal{H}}^1(X_0, B_1) = 0$ by Lemma 4.9. It follows that $\text{Ext}_{\mathcal{H}}^1(X_0, X_1) = 0$ since $\text{Ext}_{\mathcal{H}}^2(X_0, X_2) = 0$ due to $\mathcal{H}$ being hereditary. It is a contradiction. This finishes the proof for $m = 1$.

Now suppose $m > 1$. We will prove that there are at least one of complements X_j with smaller degree than m . We divide the proof into two cases: Case 1. All maps f_i

(equivalently g_i) are non-zero. Now we assume that there are no complements of X with smaller degree than m . Then by Lemma 4.7, the degrees of all X_i are m . If $d > 2$, then $\text{Ext}^2(X_0, X_2) = 0$ by Lemma 4.8, which is a contradiction to Corollary 4.5. If $d = 2$, then the same proof as above shows that $\text{Ext}^1(X_0, X_1) = 0$ which contradicts to Corollary 4.5. Therefore there is a complement of X with smaller degree than m . Case 2. There are some i such that $f_i = 0$ (equivalently $g_i = 0$). Then $X_i \cong X_{i+1}[1]$ for such i . It follows that X_{i+1} has smaller degree than X_i if X_i has strictly positive degree.

Therefore we have a complement of X , say X_s , such that the degree k'_1 of X_s is smaller than $m = k_1$. Now we replace X by $X' = (X \setminus X_0) \oplus X_s$, which is, by Theorem 4.4, a d -cluster tilting object in $\mathcal{C}_d(\mathcal{H})$, containing $|I|$ indecomposable direct summands. The number of indecomposable direct summands of X' with the (maximal) degree $m(= \nu(M))$ is $t - 1$. We repeat the step for the complement $M_2[k_2]$ of almost complete tilting object $X' \setminus M_2[k_2]$, we get a d -cluster tilting object X'' containing $|I|$ indecomposable direct summands, and the number of indecomposable direct summands of X'' with the (maximal) degree $m(= \nu(M))$ is $t - 2$. Repeating such step t times, one can get a (basic) d -cluster tilting object T containing $|I|$ indecomposable direct summands and $\nu(T) < \nu(M)$. By induction, T contains exactly n indecomposable direct summands. Then $|I| = n$. $\square$

Remark 4.10. *Theorem 4.6 is proved by Thomas in [Th] for a simply-laced Dynkin quiver (Γ, Ω_0) , there he uses the fact that $\text{ind} D^b(K\vec{\Delta}) \approx \mathbf{Z}\vec{\Delta}$ for a Dynkin quiver $\vec{\Delta}$. This fact does not hold for non-Dynkin quivers. Our proof is more categorical.*

Denote by $\mathcal{E}(\mathcal{H})$ the set of isomorphism classes of indecomposable exceptional representations in $\mathcal{H}$. The set $\mathcal{E}(\mathcal{C}_d(\mathcal{H}))$ of isoclasses of indecomposable exceptional objects in $\mathcal{C}_d(\mathcal{H})$ is the (disjoint) union of subsets $\mathcal{E}(\mathcal{H})[i]$, $i = 0, 1, \dots, d - 1$, with $\{P_j[d] \mid 1 \leq j \leq n\}$. A subset $\mathcal{M}$ of $\mathcal{E}(\mathcal{C}_d(\mathcal{H}))$ is called exceptional if for any $X, Y \in \mathcal{M}$, $\text{Ext}^i(X, Y) = 0$ for all $i = 1, \dots, d$. Denote by $\mathcal{E}_+(\mathcal{C}_d(\mathcal{H}))$ the subset of $\mathcal{E}(\mathcal{C}_d(\mathcal{H}))$ consisting of all indecomposable exceptional objects other than $P_1[d], \dots, P_n[d]$.

Now we are ready to define a simplicial complex associated to the d -cluster category $\mathcal{C}_d(\mathcal{H})$, which is a generalization of the classical cluster complexes of cluster categories [BMRRT, Rin, Z1].

Definition 4.11. *The cluster complex $\Delta^d(\mathcal{H})$ of $\mathcal{C}_d(\mathcal{H})$ is a simplicial complex which has $\mathcal{E}(\mathcal{C}_d(\mathcal{H}))$ as the set of vertices, and has exceptional subsets in $\mathcal{C}_d(\mathcal{H})$ as its simplices. The positive part $\Delta_+^d(\mathcal{H})$ is the subcomplex of $\Delta^d(\mathcal{H})$ on the subset $\mathcal{E}_+(\mathcal{C}_d(\mathcal{H}))$.*

From the definition, the facets (maximal simplices) are exactly the d -cluster tilting subsets (i.e. the sets of indecomposable objects of $\mathcal{C}_d(\mathcal{H})$ (up to isomorphism) whose direct sum is a d -cluster tilting object).

Proposition 4.12. *1. $\Delta^d(\mathcal{H})$ and $\Delta_+^d(\mathcal{H})$ are pure of dimension $n - 1$,*

- 2. For any sink (or source) k , the BGP-reflection functor $\tilde{S}_k^+$ (resp. $\tilde{S}_k^-$) induces an isomorphism between $\Delta^d(\mathcal{H})$ and $\Delta^d(s_k \mathcal{H})$. In particular, if Γ is a Dynkin diagram and Ω and Ω' are two orientations of Γ , then $\Delta^d(\mathcal{H})$ and $\Delta^d(\mathcal{H}')$ are isomorphic.*

- Proof.* 1. It follows from Theorem 4.6 that any d -cluster tilting subset contains exactly n elements. Hence $\Delta^d(\mathcal{H})$ is pure of dimension $n - 1$. Now suppose $M = \bigoplus_{i=1}^{n-1} M_i$ is an exceptional object in $\mathcal{C}_d(\mathcal{H})$ and none of the M_i are isomorphic to $P_j[d]$ for any j . In the proof of Theorem 4.6, we proved that not all complements of an almost complete tilting objects have the same degrees. Then M has a complement in $\mathcal{E}_+(\mathcal{C}_d(\mathcal{H}))$. This proves that $\Delta_+^d(\mathcal{H})$ is pure of dimension $n - 1$.
2. Since $\tilde{S}_k^+$ is a triangle equivalence from the d -cluster category $\mathcal{C}_d(\mathcal{H})$ to $\mathcal{C}_d(s_k\mathcal{H})$, it sends (indecomposable) exceptional objects to (indecomposable) exceptional objects. Thus it induces an isomorphism from $\Delta^d(\mathcal{H})$ to $\Delta^d(s_k\mathcal{H})$. The second statement follows from the first statement together with that for two orientations Ω, Ω' of a Dynkin graph Γ , there is a admissible sequence with respect to sinks $i_1, \dots, i_n$ such that $\Omega' = s_{i_n} \cdots s_{i_1} \Omega$.

□

5 Cluster combinatorics of d -cluster categories

We now define a map $\gamma_{\mathcal{H}}^d$ from $\text{ind}\mathcal{C}_d(\mathcal{H})$ to $\Phi_{\geq -1}^d$. Note that any indecomposable object X of degree i in $\mathcal{C}_d(\mathcal{H})$ has the form $M[i]$ with $M \in \text{ind}\mathcal{H}$, and if $i = d$ then $M = P_j$, an indecomposable projective representation.

Definition 5.1. Let $\gamma_{\mathcal{H}}^d$ be defined as follows. Let $M[i] \in \text{ind}\mathcal{C}_d(\mathcal{H})$, where $M \in \text{ind}\mathcal{H}$ and $i \in \{1, \dots, d\}$ (note that if $i = d$ then $M = P_j$ for some j). We set

$$\gamma_{\mathcal{H}}^d(M[i]) = \begin{cases} (\underline{\dim} M)^{i+1} & \text{if } M[i] \in \text{ind}\mathcal{H}[i], \text{ for some } 0 \leq i \leq d-1; \\ (-\alpha_j)^1 & \text{if } M[i] = P_j[d], \end{cases}$$

This map is one kind of extension of correspondence in Gabriel-Kac's Theorem between the indecomposable representations of quivers and positive roots of corresponding Lie Kac-Moody algebras. It is a bijection if Γ is a Dynkin diagram.

We denote by $\Phi_{>0}^{sr}$ the set of real Schur roots of (Γ, Ω) , i.e.

$$\Phi_{>0}^{sr} = \{ \underline{\dim} M \mid M \in \text{ind}\mathcal{E}(\mathcal{H}) \}.$$

Then the map $M \mapsto \underline{\dim} M$ gives a 1-1 correspondence between $\mathcal{E}(\mathcal{H})$ and $\Phi_{>0}^{sr}$ [Rin].

If we denote the set of colored almost positive real Schur roots by $\Phi_{\geq -1}^{sr,d}$, which consists of, by definition, d copies of the set $\Phi_{>0}^{sr}$ together with one copy of the negative simple roots, then the map $\gamma_{\mathcal{H}}^d$ gives a bijection from $\mathcal{E}(\mathcal{C}_d(\mathcal{H}))$ to $\Phi_{\geq -1}^{sr,d}$. $\Phi_{\geq -1}^{sr,d}$ contains a subset $\Phi_{>0}^{sr,d}$ consisting of all colored positive real Schur roots. The restriction of $\gamma_{\mathcal{H}}^d$ gives a bijection from $\mathcal{E}_+(\mathcal{C}_d(\mathcal{H}))$ to $\Phi_{>0}^{sr,d}$.

Since $\mathcal{E}(\mathcal{H}) \xrightarrow{\gamma_{\mathcal{H}}^{sr}} \Phi_{>0}^{sr} : M \mapsto \underline{\dim} M$ is a bijection, we use M_{β} to denote the unique indecomposable exceptional representation in $\mathcal{H}$ whose dimension vector is β . It follows from Proposition 4.2 that $\gamma_{\mathcal{H}}^d(M_{\beta}[i]) = \beta^{i+1}$ for any $0 \leq i \leq d-1$. We sometimes use

$M_{\beta^{i+1}}$ to denote the unique preimage of a colored almost positive real Schur root β^{i+1} under $\gamma_{\mathcal{H}}^d$.

We now prepare to define a simplicial complex $\Delta^{d,\mathcal{H}}(\Phi)$ associated with any root system Φ , which turns out to be isomorphic to the cluster complex $\Delta^d(\mathcal{H})$ of the d -cluster category $\mathcal{C}_d(\mathcal{H})$. When Γ is a Dynkin graph, taking an alternating orientation Ω_0 of Γ , this complex $\Delta^{d,\mathcal{H}_0}(\Phi)$ is the generalized cluster complex $\Delta^d(\Phi)$ defined by Fomin and Reading [FR2].

First of all, we define the " d -compatibility degree" on any pair of colored almost positive real Schur roots.

Definition 5.2. *For any pair of colored almost positive real Schur roots α, β , the d -compatibility degree of α, β is defined as follows:*

$$(\alpha||\beta)_{d,\mathcal{H}} = \dim_{\text{End}M_\alpha}(\text{Ext}^1(M_\alpha, \oplus_{i=0}^{i=d-1} M_\beta[i])),$$

where $\dim_{\text{End}M_\alpha}(\text{Ext}^1(M_\alpha, \oplus_{i=0}^{i=d-1} M_\beta[i]))$ denotes the length of $\text{Ext}^1(M_\alpha, \oplus_{i=0}^{i=d-1} M_\beta[i])$ as a right $\text{End}M_\alpha$ -module. When $d > 1$, $\text{End}M_\alpha$ is a division algebra by Proposition 4.2 (3), and this length equals the dimension of $\text{Ext}^1(M_\alpha, \oplus_{i=0}^{i=d-1} M_\beta[i])$ over the division algebra $\text{End}M_\alpha$.

Remark 5.3. *When Γ is a Dynkin diagram with trivial valuation and Ω_0 is an alternating orientation of Γ , this compatibility degree is defined in [Th]. When $d = 1$ and Γ is a Dynkin diagram, we recover the classical compatibility degree defined in [BMRRT, Z2].*

Theorem 5.4. 1. *For any pair of colored almost positive real Schur roots α, β ,*

- (a) $(\alpha||\beta)_{d,\mathcal{H}} = (\sigma_{k,d}(\alpha)||\sigma_{k,d}(\beta))_{d,s_k\mathcal{H}}$, if k is a sink (or a source);
- (b) $(\alpha||\beta)_{d,\mathcal{H}} = (R_{d,\Omega}(\alpha)||R_{d,\Omega}(\beta))_{d,\mathcal{H}}$,
- (c) $(\alpha||\beta)_{d,\mathcal{H}} = 0$ if and only if $(\beta||\alpha)_{d,\mathcal{H}} = 0$.

- 2. *For any almost positive real Schur root β , $((-\alpha_i)^1||(\beta)^l)_{d,\mathcal{H}} = 0$ if and only if $\max\{n_i(\beta), 0\} = 0$ where $n_i(\beta)$ is the coefficient of α_i in the expansion of β in terms of the simple roots $\alpha_1, \dots, \alpha_n$.*

Proof. 1. Let α, β be two colored almost positive real Schur roots.

- (a) We prove it for the case k is a sink, the proof for source is similar. It is easy to check that the following diagram is commutative:

$$\begin{array}{ccc} \text{ind}\mathcal{C}_d(\mathcal{H}) & \xrightarrow{\tilde{S}_k^+} & \text{ind}\mathcal{C}_d(s_k\mathcal{H}) \\ \gamma_{\mathcal{H}}^d \downarrow & & \downarrow \gamma_{s_k\mathcal{H}}^d \\ \Phi_{\geq -1}^d & \xrightarrow{\sigma_{k,d}} & \Phi_{\geq -1}^d \end{array}$$

Hence we have that

$$\begin{aligned} (\sigma_{k,d}(\alpha)||\sigma_{k,d}(\beta))_{d,s_k\mathcal{H}} &= \dim_{\text{End}\tilde{S}_k^+(M_\alpha)} \text{Ext}^1(\tilde{S}_k^+(M_\alpha), \oplus_{i=0}^{i=d-1} \tilde{S}_k^+(M_\beta)[i]) \\ &= \dim_{\text{End}M_\alpha} \text{Ext}^1(M_\alpha, \oplus_{i=0}^{i=d-1} M_\beta[i]) \\ &= (\alpha||\beta)_{d,\mathcal{H}} \end{aligned}$$

- (b) As we mentioned before, the shift functor $[1]$ of $\mathcal{C}_d(\mathcal{H})$ is an auto-equivalence. We now check the following diagram commutes:

$$\begin{array}{ccc} \mathrm{ind}\mathcal{C}_d(\mathcal{H}) & \xrightarrow{[1]} & \mathrm{ind}\mathcal{C}_d(\mathcal{H}) \\ \gamma_{\mathcal{H}}^d \downarrow & & \downarrow \gamma_{\mathcal{H}}^d \\ \Phi_{\geq -1}^d & \xrightarrow{R_{d,\Omega}} & \Phi_{\geq -1}^d \end{array}$$

By Proposition 2.2, any indecomposable object in $\mathcal{C}_d(\mathcal{H})$ is of the form $X[i]$ with X an indecomposable representation in $\mathcal{H}$ and with $0 \leq i \leq d-1$, or of the form $P_j[d]$. Denote by $\underline{\dim} X = \alpha$. If $i \leq d-2$, then $R_{d,\Omega}\gamma_{\mathcal{H}}^d(X[i]) = R_{d,\Omega}((\alpha)^{i+1}) = (\alpha)^{i+2} = \gamma_{\mathcal{H}}^d[1](X[i])$. We will prove the equality for other indecomposable objects in $\mathcal{C}_d(\mathcal{H})$. Firstly we have that $R_{d,\Omega}\gamma_{\mathcal{H}}^d(P_j[d-1]) = R_{d,\Omega}((\underline{\dim} P_j)^d) = (-\alpha_j)^1$ and $\gamma_{\mathcal{H}}^d[1](P_j[d-1]) = (-\alpha_j)^1$. Hence $R_{d,\Omega}\gamma_{\mathcal{H}}^d(P_j[d-1]) = \gamma_{\mathcal{H}}^d[1](P_j[d-1])$. Secondly, for any $X[d-1]$ with X not being projective, we have $\tau X \in \mathrm{ind}\mathcal{H}$. We have that $R_{d,\Omega}\gamma_{\mathcal{H}}^d(X[d-1]) = R_{d,\Omega}((\alpha)^d) = (R_{\Omega}(\alpha))^1$ and $\gamma_{\mathcal{H}}^d[1](X[d-1]) = \gamma_{\mathcal{H}}^d(X[d]) = \gamma_{\mathcal{H}}^d(\tau^{-1}[d]\tau X) = \gamma_{\mathcal{H}}^d(\tau X) = (R_{\Omega}(\alpha))^1$. The last equality holds since $\underline{\dim} \tau X = R_{\Omega}(\underline{\dim} X)$ (compare Section 3.1). This proves that $R_{d,\Omega}\gamma_{\mathcal{H}}^d(X[d-1]) = \gamma_{\mathcal{H}}^d[1](X[d-1])$. Similar proof for $P_j[d]$. We finish the proof of the commutativity of the diagram.

It follows that

$$\begin{aligned} (R_{d,\Omega}(\alpha) || R_{d,\Omega}(\beta))_{d,\mathcal{H}} &= \dim_{\mathrm{End}_{M_{\alpha}[1]}} \mathrm{Ext}^1(M_{\alpha}[1], \oplus_{i=0}^{i=d-1} M_{\beta}[1][i]) \\ &= \dim_{\mathrm{End}_{M_{\alpha}}} \mathrm{Ext}^1(M_{\alpha}, \oplus_{i=0}^{i=d-1} M_{\beta}[i]) \\ &= (\alpha || \beta)_{d,\mathcal{H}}, \end{aligned}$$

where the second equality follows from the fact that $[1]$ is an equivalence.

- (c) Let $X, Y \in \mathcal{C}_d(\mathcal{H})$ with $\mathrm{Ext}^i(X, Y) = 0$ for any $1 \leq i \leq d$. Then by the Calabi-Yau property of $\mathcal{C}_d(\mathcal{H})$, we have that for any $1 \leq j \leq d$ $\mathrm{Ext}^j(Y, X) \cong \mathrm{Ext}^{d-j+1}(X, Y) = 0$. This proves (c).
2. We first prove the necessity: Let β be an almost positive real Schur root with $((-\alpha_i)^1 || (\beta)^l)_{d,\mathcal{H}} = 0$. If β is a negative simple root and $l = 1$, we have easily that $\max\{n_i(\beta), 0\} = 0$. Now we assume that β is a positive real Schur root. From the condition $((-\alpha_i)^1 || (\beta)^l)_{d,\mathcal{H}} = 0$, we have $\mathrm{Ext}^j(P_i[d], M_{\beta^l}) = 0$, i.e. $\mathrm{Ext}^j(P_i[d], M_{\beta}[l-1]) = 0$, for any $1 \leq j \leq d$. Since $1 \leq l \leq d$, we have $1 \leq j \leq d$, where $j = d+1-l$. Now we have that $0 = \mathrm{Ext}^j(P_i[d], M_{\beta}[l-1]) \cong \mathrm{Hom}(P_i[d], M_{\beta}[l+j-1]) \cong \mathrm{Hom}(P_i, M_{\beta})$. Hence $n_i(\beta) = \dim_{\mathrm{End} P_i} \mathrm{Hom}(P_i, M_{\beta}) = 0$.

Now we prove the other direction. Suppose that β is an almost positive real Schur root with $\max\{0, n_i(\beta)\} = 0$. Firstly, if β is the negative of a simple root, say $(-\alpha_j)^1$, then

$$\begin{aligned} ((-\alpha_i)^1 || (-\alpha_j)^1)_{d,\mathcal{H}} &= \dim_{\mathrm{End}(P_i[d])} \mathrm{Ext}^1(P_i[d], \oplus_{k=0}^{k=d-1} P_j[d][k]) \\ &= \dim_{\mathrm{End}(P_i[d])} \mathrm{Ext}^1(P_i, \oplus_{k=0}^{k=d-1} P_j[k]) \\ &= \dim_{\mathrm{End}(P_i[d])} \mathrm{Ext}^1(P_i, P_j[d-1]), \end{aligned}$$

the last equality following from Proposition 4.2 (4). But $\text{Ext}^1(P_i, P_j[d-1]) \cong \text{Hom}(P_i, P_j[-1][d+1]) \cong D\text{Hom}(P_j[-1], P_i) \cong D\text{Ext}^1(P_j, P_i) = 0$. This proves that $((-\alpha_i)^1 || (-\alpha_j)^1)_{d, \mathcal{H}} = 0$. Now we assume that β is a positive real Schur root and l is a positive integer not exceeding d . We will prove that $((-\alpha_i)^1 || (\beta)^l)_{d, \mathcal{H}} = 0$ under the condition that $n_i(\beta) = 0$. We can assume that $d > 1$ since for $d = 1$, the corresponding result is proved in [Z1]. It follows from the condition $n_i(\beta) = 0$ that $\text{Hom}_{\mathcal{H}}(P_i, M_\beta) = 0$, and then $\text{Hom}(P_i, M_\beta) = 0$. Hence $\text{Ext}^1(P_i[d], M_\beta[d-1]) \cong \text{Hom}(P_i, M_\beta) = 0$. We will prove that $\text{Ext}^j(P_i[d], M_{\beta^l}) = 0$ for $1 \leq j \leq d$. Now given such j , $\text{Ext}^j(P_i[d], M_{\beta^l}) = \text{Ext}^j(P_i[d], M_\beta[l-1]) = \text{Ext}^1(P_i[d], M_\beta[l+j-2]) \cong \text{Ext}^1(P_i, M_\beta[l+j-d-2])$. Since $1 \leq l \leq d$, $1 \leq j \leq d$, we have $-d \leq l+j-d-2 \leq d-2$. Then we have that $\text{Ext}^j(P_i[d], M_{\beta^l}) = 0$ which follows from Proposition 4.2 (4) for $l+j-d-2 \neq -1$ and from the fact that $\text{Ext}^1(P_i, M_\beta[-1]) \cong \text{Hom}(P_i, M_\beta) = 0$ for $l+j-d-2 = -1$. $\square$

Definition 5.5. Let Φ be the root system corresponding to Γ and $\mathcal{H}$ the category of representations of the valued quiver (Γ, Ω) .

1. Any pair α, β of almost positive real Schur roots is called d -compatible if $(\alpha || \beta)_{d, \mathcal{H}} = 0$; a subset of $\Phi_{\geq -1}^{sr, d}$ is called d -compatible if any two elements of this subset are compatible.
2. The simplicial complex $\Delta^{d, \mathcal{H}}(\Phi)$ associated to Φ and $\mathcal{H}$ is a complex which has $\Phi_{\geq -1}^{sr, d}$ as the set of vertices. Its simplices are d -compatible subsets of $\Phi_{\geq -1}^{sr, d}$. The subcomplex of $\Delta^{d, \mathcal{H}}(\Phi)$ which has $\Phi_{\geq 0}^{sr, d}$ as the set of vertices is denoted by $\Delta_+^{d, \mathcal{H}}(\Phi)$. We call $\Delta^{d, \mathcal{H}}(\Phi)$ the generalized cluster complex associated to Φ and $\mathcal{H}$.

Remark 5.6. Given a graph Γ , we have the corresponding root system Φ . Since the set of real Schur roots of Φ depends on the category $\text{ind} \mathcal{H}$, equivalently, on the orientation Ω of Γ , the generalized cluster complexes $\Delta^{d, \mathcal{H}}(\Phi)$ are possibly non-isomorphic for different orientations of Γ , but they are isomorphic to each other if Γ is a Dynkin diagram by Proposition 4.12 (2) and the following theorem.

Theorem 5.7. 1. Let Γ be a valued graph and Φ the corresponding root system. Let Ω be an admissible orientation of Γ . Then $\gamma_{\mathcal{H}}^d$ provides an isomorphism from the simplicial complex $\Delta^d(\mathcal{H})$ to the generalized cluster complex $\Delta^{d, \mathcal{H}}(\Phi)$, which sends vertices to vertices, k -faces to k -faces.

2. The restriction of $\gamma_{\mathcal{H}}^d$ to $\Delta_+^d(\mathcal{H})$ gives an isomorphism from $\Delta_+^d(\mathcal{H})$ to $\Delta_+^{d, \mathcal{H}}(\Phi)$.
3. If Γ is a Dynkin graph and Ω_0 is an alternating orientation of Γ , then $\Delta^{d, \mathcal{H}_0}(\Phi)$ is the generalized cluster complex $\Delta^d(\Phi)$ defined by Fomin-Reading in [FR2].

Proof. 1. $\gamma_{\mathcal{H}}^d$ provides a bijection from the vertices of $\Delta^d(\mathcal{H})$ to that of $\Delta^{d, \mathcal{H}}(\Phi)$. For any pair of colored almost positive real Schur roots α^k, β^l , they are d -compatible if and only if $M_{\alpha^k} \oplus M_{\beta^l}$ is an exceptional object where M_{α^k} and M_{β^l} are the exceptional objects corresponding to α^k, β^l respectively under the map $\gamma_{\mathcal{H}}^d$. Hence $\gamma_{\mathcal{H}}^d$ is an isomorphism from $\Delta^d(\mathcal{H})$ to $\Delta^{d, \mathcal{H}}(\Phi)$.

2. This is a direct consequence of 1.
3. This is a direct consequence of Theorem 3.9 and Theorem 5.4.

□

From Theorem 5.7, one can translate results from each side. For example, one gets the number of d -cluster tilting objects in $\mathcal{C}_d(\mathcal{H})$ from the number of facets of generalized cluster complexes of finite root systems [FR2].

Corollary 5.8. 1. *The generalized cluster complex $\Delta^{d,\mathcal{H}}(\Phi)$ and its subcomplex $\Delta_+^{d,\mathcal{H}}(\Phi)$ are pure of dimension $n - 1$.*

2. *Let (Γ, Ω) be a connected Dynkin quiver and Φ the root system corresponding to Γ . Then the number of d -cluster tilting objects of $\mathcal{C}_d(\mathcal{H})$ is $\prod_i \frac{dh+e_i+1}{e_i+1}$, where h is the Coxeter number of Φ and $e_1, \dots, e_n$ the exponents of Φ .*
3. *Let (Γ, Ω) be a connected Dynkin quiver and Φ the corresponding root system. Then the number of complements of any almost complete tilting object in $\mathcal{C}_d(\mathcal{H})$ is $d + 1$.*

Proof. 1. It follows from Proposition 4.12(1) and Theorem 5.7(1).

2. It follows from Theorem 5.7(1) and Proposition 8.4 in [FR2] that the statement in 2. holds for the d -cluster category $\mathcal{C}_d(\mathcal{H}_0)$ of Ω_0 . Then by Proposition 4.12(2), the statement in 2. holds for a d -cluster category $\mathcal{C}_d(\mathcal{H})$ corresponding to an arbitrary orientation Ω .
3. It follows from Theorem 5.7(1) and Proposition 3.10 in [FR2] that the number of complements of any almost complete tilting object in $\mathcal{C}_d(\mathcal{H}_0)$ is $d + 1$. Hence by Proposition 4.12(2) the number of complements of any almost complete tilting object in $\mathcal{C}_d(\mathcal{H})$ is $d + 1$.

□

Remark 5.9. *Corollary 5.8 (1) generalizes Theorem 2.9 in [FR2] to infinite root systems.*

Remark 5.10. 1. *From Corollary 5.8 (2) for $d = 1$, combining with the result in [BMRRT] (see also [KZ]), in which the cluster tilting subcategories in $\mathcal{D}$ are proved to be in one-to-one correspondence with the cluster tilting modules in cluster categories by the projection π , we have an explanation on why the number of cluster tilting subcategories (i.e. Ext-configurations in [I1]) in $\mathcal{D}$ is the same as the number of facets of $\Delta(\Phi)$.*

2. *Corollary 5.7(3) is proved by Thomas in [Th] for an alternating simply-laced Dynkin quiver (Γ, Ω_0) , using a different approach.*

ACKNOWLEDGMENTS.

The author would like to thank Claus Michael Ringel and Osamu Iyama for his helpful suggestions on cluster complexes and the formalism of the paper and to thank Bernhard Keller for informing the author of Y. Palu's work. After completing this work, I was told by Bernhard Keller and Robert Marsh that H.Thomas had some similar results for simply-laced Dynkin type [Th]. I thank them very much for this!

The author would like to thank the referees for their useful suggestions to improve the paper.

References

- [ABST] I. Assem, T. Brstle, R. Schiffler and G. Todorov, m -cluster categories and m -replicated algebras. Preprint [arXiv:math.RT/0608727](#), 2006.
- [AT] C. Athanasiadis and E. Tzanaki. Shellability and higher Cohen-Macaulay connectivity of generalized cluster complexes. Preprint [arXiv:math.CO/0606018](#), 2006.
- [BaM] K.Baur and R.Marsh. A Geometric description of m -Cluster categories. Preprint, [math.RT/0610512](#). To appear in Transactions of the AMS.
- [BM] A.Buan and R.Marsh. Cluster-tilting theory. Trends in representation theory of algebras and related topics, Edited by J.de la Peña and R. Bautista. Contemporary Mathematics. **406**, 1-30, 2006.
- [BMR] A.Buan, R.Marsh, and I.Reiten. Cluster mutation via quiver representations. Preprint, [arXiv:math.RT/0412077](#), 2004. To appear in Comment. Math.Helv.
- [BMRRT] A.Buan, R.Marsh, M.Reineke, I.Reiten and G.Todorov. Tilting theory and cluster combinatorics. Advances in Math. **204**, 572-618, 2006.
- [CC] P.Caldero and F.Chapoton. Cluster algebras as Hall algebras of quiver representations. Comment. Math. Helv. **81**, 595-616, 2006
- [CCS] P.Caldero, F.Chapoton and R.Schiffler. Quivers with relations arising from clusters (A_n case). Transactions of the AMS. **358**, 1347-1364, 2006.
- [CK] P.Caldero and B.Keller. From triangulated categories to cluster algebras. To appear in Invent.Math. [arXiv:math.RT/0506018](#).
- [DR] V. Dlab and C. M. Ringel. Indecomposable representations of graphs and algebras. Mem. Amer. Math. Soc. **591** (1976).
- [FR1] S.Fomin and N.Reading. Root system and generalized associahedra, lecture notes for the IAS/Park City Graduate Summer School in Geometric Combinatorics, 2004.
- [FR2] S.Fomin and N.Reading. Generalized cluster complexes and Coxeter combinatorics. IMRN. **44**, 2709-2757, 2005.
- [FZ1] S.Fomin and A.Zelevinsky. Cluster Algebras I: Foundations. Jour. Amer. Math. Soc. **15**, no. 2, 497-529, 2002.
- [FZ2] S.Fomin and A.Zelevinsky. Y-system and generalized associahedra, Ann of Math. **158**, 977-1018, 2003.
- [FZ3] S.Fomin and A.Zelevinsky. Cluster algebras II: Finite type classification. Invent. Math. **154**, no.1, 63-121, 2003.
- [I1] O.Iyama. Higher dimensional Auslander-Reiten theory on maximal orthogonal subcategories. To appear in Adv. Math. See also [arXiv:math.RT/0407052](#).

- [I2] O.Iyama. Maximal orthogonal subcategories of triangulated categories satisfying Serre duality, Mathematisches Forschungsinstitut Oberwolfach Report, no. 6(2005), 353-355.
- [IY] O.Iyama and Y. Yoshino. Mutations in triangulated categories and rigid Cohen-Macaulay modules, Preprint. [arXiv:math.RT/0607736](#).
- [Ke] B.Keller. Triangulated orbit categories. Documenta Math. **10**, 551-581, 2005.
- [KR] B.Keller and I.Reiten. Cluster-tilted algebras are Gorenstein and stably Calabi-Yau. To appear in Adv. Math. See also [arXiv:math.RT/0512471](#).
- [KZ] S.Koenig, and B.Zhu. From triangulated categories to abelian categories– cluster tilting in a general framework. To appear in Math. Zeit. See also [arXiv:math.RT/0605100](#).
- [MRZ] R. Marsh, M. Reineke and A. Zelevinsky. Generalized associahedra via quiver representations. Transactions of AMS. **355(10)**(2003), PP. 4171-4186.
- [Pa] Y. Palu. Ph.D Thesis in preparation.
- [Rin] C. M. Ringel. Some remarks concerning tilting modules and tilted algebras. Origin. Relevance. Future. An appendix to the Handbook of tilting theory, edited by Lidia Angeleri-Hügel, Dieter Happel and Henning Krause. Cambridge University Press (2007), LMS Lecture Notes Series 332.
- [Th] H.Thomas. Defining an m -cluster category. , Preprint, 2005.
- [Z1] B.Zhu. BGP-reflection functors and cluster combinatorics, Jour. of Pure and Applied Algebra **209**, 497-506, 2007.
- [Z2] B.Zhu. Equivalences between cluster categories, Jour. of Algebra. **304**(2006), 832-850.